



# Kent Academic Repository

**Watkins, Richard, Littlefair, P, Kolokotroni, M and Palmer, J (2002) *The London Heat Island – surface and air temperature measurements in a park and street gorges*. ASHRAE Transactions, 108 (1). pp. 419-427. ISSN 0001-2505.**

## Downloaded from

<https://kar.kent.ac.uk/52517/> The University of Kent's Academic Repository KAR

## The version of record is available from

## This document version

UNSPECIFIED

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

# The London Heat Island—Surface and Air Temperature Measurements in a Park and Street Gorges

Richard Watkins

John Palmer, C.Eng.

Maria Kolokotroni, Ph.D.  
Member ASHRAE

Paul Littlefair, Ph.D., C.Eng.

## ABSTRACT

*This paper reports results from short-term tests carried out as part of a project to characterize the urban heat island in London. The investigations looked at air temperatures upstream and downstream of a park and the surface and air temperatures within street gorges. It was found that the air in the park was associated with lower mean ( $0.6^{\circ}\text{C}$  [ $1.1^{\circ}\text{F}$ ] less) and peak temperatures ( $1.1^{\circ}\text{C}$  [ $2.0^{\circ}\text{F}$ ] less) compared to residential or shopping streets on either side. The apparent cooling influence of the park extended downstream between 200 and 400 meters (200 to 400 yards). Measurements in four street gorges showed a wide variation in surface temperatures—up to  $22^{\circ}\text{C}$  ( $40^{\circ}\text{F}$ )—although  $5^{\circ}\text{C}$  to  $10^{\circ}\text{C}$  ( $9^{\circ}\text{F}$  to  $18^{\circ}\text{F}$ ) was more typical. For a given facade, lighter surfaces were associated with lower temperatures, between  $6^{\circ}\text{C}$  and  $10^{\circ}\text{C}$  ( $11^{\circ}\text{F}$  and  $18^{\circ}\text{F}$ ) cooler. A strong relationship was found between mean gorge surface temperature and the gorge air temperature measured at 6 m (20 ft) (half-gorge height). This was true for both a sunny day and a cloudy day. The results suggest that significant reductions in air temperature may be possible by adjusting the albedo of urban surfaces.*

## INTRODUCTION

The heat island effect leads to urban air temperatures being, in general, warmer than those found in rural areas. The form and fabric of an urban area differ substantially from rural zones, and this alters the way in which heat flows into and out of the area. Solar gain is more efficiently captured in a city and, together with heat from anthropogenic sources, is better stored. Likewise, the way in which heat may be dispersed through re-radiation and air cooling is less effective than in the countryside. The effect of these urban/rural differences can be

seen if temperature is measured along a transect from the countryside into a city. However, a city is a complex of microclimates with local variation dependent on form, surface characteristics, and anthropogenic fluxes. This variation can act as a guide to developing ways of modifying urban temperatures and mitigating the heat island effect.

The heat island in London is being monitored each hour at 80 measurement stations located on a uniform radial grid of approximate diameter 50 km (30 miles) and some of the results have been reported by the authors elsewhere (Palmer et al. 2000; Watkins et al. 2001). To supplement the long-term air temperature data, short-term tests were carried out in the summer of 2000, aimed at identifying the effect of a park on air temperature and the relationship between air and surface temperature within street gorges on sunny and cloudy days. These measurements are reported in this paper.

## Parks

Parks usually contain vegetation through which evapotranspiration converts incoming solar radiation into latent rather than sensible heat, thereby reducing the surface temperature and, in turn, the air temperature. Elevated vegetation, such as tree canopies, intercepts solar energy before it reaches the ground, both providing shading and a better opportunity for the heat to be dispersed. Many observations have confirmed the cooling effect of parks, although their influence beyond their boundaries is uncertain.

Landsberg (1981) reported that large parks situated in urban areas are cooler on warm summer nights. Saito has investigated air and surface temperatures in Kumamoto City in Japan and found that the urban air temperature distribution is related to the distribution of green covering (Saito et al.

---

Richard Watkins is a research student and Maria Kolokotroni is a lecturer at Brunel University, Uxbridge, Middlesex, U.K. John Palmer and Paul Littlefair are principal consultants at Building Research Establishment, Garston, Watford, U.K.